

ADMIT to Infusion Center of EAMC for Eculizumab (Soliris)

Diagnosis:

- | | |
|---|--|
| <input type="checkbox"/> Atypical Hemolytic Uremic Syndrome | <input type="checkbox"/> Generalized Myasthenia Gravis, Refractory |
| <input type="checkbox"/> Neuromyelitis Optica Spectrum Disorder | <input type="checkbox"/> Paroxysmal Nocturnal Hemoglobinuria |
| <input type="checkbox"/> Other (specify) _____ | |

Labs:

- Labs must be resulted prior to scheduling infusion.
- Physician office to verify that patient has received a meningococcal vaccine at least 14 days prior to therapy and the patient has been revaccinated according to current guidelines.
- Physician office to verify patient is enrolled in the REMS program.

Pre-Meds:

- | | |
|---|--|
| <input type="checkbox"/> Acetaminophen 650 mg PO with each treatment | <input type="checkbox"/> Methylprednisolone 125 mg IVP with each treatment |
| <input type="checkbox"/> Diphenhydramine 25 mg PO with each treatment | <input type="checkbox"/> Famotidine 20 mg IVP over 2 minutes with each treatment |
| <input type="checkbox"/> Diphenhydramine 50 mg PO with each treatment | <input type="checkbox"/> Other: _____ |

Infusion Orders

Drug	Fluid	Dose	Route
Eculizumab (Soliris)	Normal Saline (see below)*	<input type="checkbox"/> 600 mg <input type="checkbox"/> 900 mg <input type="checkbox"/> 1,200 mg	IV Infusion

Induction Dose:

- ☐ 600 mg weekly for 4 doses
- ☐ 900 mg weekly for 4 doses

Maintenance Dose:

- ☐ 900 mg at week 5; then 900 mg every 2 weeks thereafter.
- ☐ 1,200 mg at week 5; then 1,200 mg every 2 weeks thereafter.

Ancillary Treatment/Administrative Procedures

- ❖ *Eculizumab should be diluted with an equal volume of Normal Saline to a final concentration of 5 mg/ml.
- ❖ Infuse over 35 minutes. Decrease infusion rate or discontinue for infusion reactions; do not exceed a maximum 2-hour duration of infusion.
- ❖ Monitor vital signs prior to infusion, then as needed. Monitor for at least 1 hour following completion of infusion for signs/symptoms of infusion reaction.
- ❖ May use Infusion Center at EAMC Non-Oncology standing orders.

Call 334-528-6320 and notify Neurology Center of East Alabama office of the next infusion date scheduled.

Physician Signature

Date

Time

**EAST ALABAMA MEDICAL CENTER
EAMC Infusion Order Set
Eculizumab (Soliris)**

Patient Identification