

ADMIT to COVID Infusion Center (CIC)/Infusion Center of EAMC for Bamlanivimab

Diagnosis:

☐ Mild to Moderate COVID-19

Pre-meds:

☐ Acetaminophen 650mg PO Once

☐ Diphenhydramine 25mg PO Once

☐ Diphenhydramine 50mg PO Once

☐ Other: _____

Infusion Orders

Drug	Fluid	Dose	Route
Bamlanivimab	NS 50 ml	700 mg	IV Infusion

Ancillary Treatment/Administrative Procedures

- ❖ Infuse over 16 minutes.
- ❖ Use 0.2-micron filter set.
- ❖ Monitor vital signs prior to start of infusion and for at least one hour after infusion is complete.
- ❖ Monitor for signs/symptoms of a clinically significant hypersensitivity reaction, including anaphylaxis (fever, chills, nausea, headache, bronchospasm, hypotension, angioedema, throat irritation, rash including urticaria, pruritis, myalgia, dizziness, etc.)
- ❖ May use Infusion Center at EAMC Non-Oncology standing orders.
- ❖ If an infusion reaction occurs, immediately discontinue infusion and administer medications and/or supportive care per Biologics reaction on standing orders.
- ❖ Once infusion is complete, flush the infusion line with 10 ml of 0.9% Normal Saline to ensure delivery of the required dose.

Ordering Physician

Date

Time

**EAST ALABAMA MEDICAL CENTER
Infusion Order Set
Bamlanivimab**

Patient Identification