

DATE & TIME	PHYSICIAN STANDING ORDERS for INFUSION CENTER at EAMC Non-Oncology
	BIOLOGICS REACTIONS: • STOP Infusion. • Methylprednisolone sodium succinate (SoluMedrol) 125 mg IV push x 1 • Famotidine (Pepcid) 20 mg IVP over 2 minutes with each treatment • Diphenhydramine (Benadryl) 50 mg IV slow push x 1 • 0.9% NS IV bolus up to 500 ml x 1 • Notify Physician
	PORT WITHOUT BLOOD RETURN: • Instill Alteplase 2 mg to port per policy “Dec clotting of central venous catheter” • If no results, obtain order for portogram.
	DYSPNEA: • O₂ at 2 liters/min. via nasal cannula or mask. May titrate up to 10 liters/min if dyspnea persists. • Notify Physician if dyspnea persists
	PORTACATH ACCESS: Lidocaine & Prilocaine Cream packet. Apply to skin over Portacath 30 minutes prior to access x 1.
	NAUSEA/VOMITING: Ondansetron (Zofran) 4mg IV push x 1.
	NEUROLOGY CENTER OF EAST ALABAMA PATIENTS: Ocrevus reaction or Tysabri reactions: • First reaction: • STOP infusion. • Methylprednisolone sodium succinate (SoluMedrol) 125 mg IV push x 1 • Diphenhydramine (Benadryl) 50 mg IV slow push x 1 • Second Reaction: • STOP infusion • Famotidine (Pepcid) 20 mg IVP over 2 minutes with each treatment • Methylprednisolone sodium succinate (SoluMedrol) 125 mg IV push x 1 • Notify Physician
	DR MASSEY’S PATIENTS: Infusion Reactions: • STOP Infusion. • Hydrocortisone (Solucortef) 100 mg IV push x 1 • Famotidine (Pepcid) 20 mg IVP over 2 minutes x 1 • Notify Physician
	Dalbavancin (DALVANCE) REACTIONS: • Moderate: Patient with Flushing or Pruritus but hemodynamically stable and not experiencing chest pain or muscle spasm: • Diphenhydramine (Benadryl) 50 mg IV x1 • Famotidine (Pepcid) 20 mg IVP over 2 minutes x 1 • Slow Dalvance infusion time to 1 hour • Notify Physician • Pretreat next scheduled dose with Diphenhydramine 50 mg IV x 1 and Pepcid 20 mg IVP over 2 minutes x 1, 30 minutes prior to Dalvance infusion and then infuse over 1 hour. • Severe Reaction: Patient is experiencing chest pain, muscle spasm or hypotension: • STOP Infusion • Diphenhydramine (Benadryl) 50 mg IV x1 • Famotidine (Pepcid) 20 mg IVP over 2 minutes x 1 • Notify Physician
	Physician Signatures on file. Dr. Christopher Adams, Dr. Zarmeena Ali, Dr. Jose Leon, Dr. Amanda Reimer, Dr. Ricardo Maldonado, Dr. Adahli Massey, Dr. Eric Hemberg